S5 Table. Results of refeeding experiments (early-starved). Initial weight at 5th instar, pupal weight in weight-gain and weight-loss groups, and pupal duration in *P. hilaris*, larvae of which were fed for 4 days prior to starvation

Regimen *	Initial weight at 5th instar (mg) §	Pupal weight in weight-gain group (mg) §	Pupal weight in weight-loss group (mg) §	Pupal duration (days) §
	Mean (S.D.) n	Mean (S.D.) n	Mean (S.D.) n	Mean (S.D.) n
4F-2S-F	351.8 <sup>a</sup> (102.0) 10	389.3° (147.9) 10	- ( - ) 0	11.7 <sup>a</sup> ( 0.8 ) 10
4F-4S-F	358.0 <sup>a</sup> ( - ) 1	454.0 <sup>a</sup> ( - ) 1	- ( - ) 0	$13.0^{a}(-)$
4F-5S-F	306.5 <sup>a</sup> (29.0) 2	374.0 <sup>a</sup> ( 91.9) 2	201.0 <sup>a</sup> ( - ) 1	$11.3^{a}(1.2)$ 3
4F-6S-F	373.1 <sup>a</sup> ( 56.7) 7	- ( - ) 0	303.0 <sup>a</sup> (53.0) 7	11.4 <sup>a</sup> ( 0.5 ) 7

<sup>\*</sup> See the footnote to S2 Table.

<sup>§</sup> Means in the same column with the same letter are not significantly different (Tukey test, p < 0.05).